

Original Research Article

Prevalence and determinants of nocturnal enuresis in school going children in Southern Maharashtra, India

Dayanand P. Nakate¹, Sunil S. Vaidya^{2*}, Shahaji Y. Gaikwad¹,
Ramesh S. Patil², Manoj S. Ghogare¹

¹Department of Paediatrics, ²Department of PSM, Ashwini Rural Medical College, Hospital and Research Centre, Kumbhari, Solapur, Maharashtra, India

Received: 14 January 2019

Accepted: 22 January 2019

*Correspondence:

Dr. Sunil S. Vaidya,

E-mail: drsvaidya@yahoo.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Nocturnal enuresis or bedwetting is a common health problem in children in the age group of 5 to 15 years. In India the prevalence of nocturnal enuresis ranges from approximately 7% to 12.6%. The prevalence from other parts of the world ranges from 6% to 42%. This study was planned to find the prevalence and determinants of nocturnal enuresis in school going children.

Methods: This cross-sectional study was conducted in four primary and secondary schools in Solapur. A predefined questionnaire, prepared in English and translated to local languages, Marathi and Kannada, was handed over to all the children attending the selected schools. Children were asked to hand over the questionnaire to their parents and return with the filled questionnaire. The questionnaire was prepared to obtain the information related to the socio-demographic factors of the family and child, frequency of enuresis and information related the risk factors. Nocturnal enuresis was defined if the frequency of enuresis was more than twice per week.

Results: The overall prevalence of nocturnal enuresis was 11.4 %. The maximum prevalence was found in the age group of 8 -9 years (22.96 %). The prevalence of nocturnal enuresis in boys (14.34 %) was significantly more than the prevalence in girls (8.31 %). The study showed that nocturnal enuresis was significantly associated with stress, poor school performance, sleep pattern (hard to awaken), family history, burning micturition etc. But it was not associated with maternal or paternal education, birth order, type of family.

Conclusions: The prevalence of nocturnal enuresis is 11.40 % in school going children and it is associated with age, sex, stress, family history, burning micturition, hyperactive child, poor school performance, sleep pattern (hard to awake) avoidance of going to micturate before sleep etc.

Keywords: Burning micturition, Nocturnal enuresis, School going children, Sleep pattern

INTRODUCTION

Nocturnal enuresis or bedwetting is a common health problem in children in the age group of 5 to 15 years. It is defined as an involuntary voiding of urine during sleep with a frequency of at least twice a week in children aged >5 years in the absence of congenital or acquired defects of the central nervous system.¹ The available data from

India shows that the prevalence of nocturnal enuresis ranges from approximately 7% to 12.6%.²⁻⁵ The prevalence from other parts of the world ranges from 6 % to 42 %.⁵⁻¹³

Enuresis can be nocturnal or daytime enuresis. It can also be primary or secondary. There are many possible causes of nocturnal enuresis including anxiety, hyperactivity

disorder, a gene located on chromosome 12q, overactive bladder, urinary tract infection, antidiuretic hormone deficiency, diabetes, small bladder, constipation etc.¹⁴ The data also suggest that the annual spontaneous resolution rate is approximately 15 to 16%.¹⁵⁻¹⁶

Nocturnal enuresis is a common problem which causes psychological trauma to the child as well as to the family which can result in social avoidance, feeling of guilt and shame, loss of self-esteem and can result in overall psychological development of the child.

There are numerous factors associated with the nocturnal enuresis. Few of the associated factors are sex, stress, parental education, school performance, sleep pattern etc.¹¹⁻¹²

The studies on prevalence and risk factors for nocturnal enuresis not conducted in rural part of Maharashtra. So, to fill of this gap in available literature, this study was planned to find the prevalence and determinants of nocturnal enuresis in school going children.

METHODS

This cross-sectional study was conducted by department of Paediatrics, Ashiwani Rural Medical College, Hospital and Research Centre, Solapur. The study period was July 2018 to August 2018. The study was conducted in four primary and secondary schools in Solapur. Two of them were Marathi medium schools and two were English medium schools.

Inclusion criteria

- School going child in the age group of 5 to 15 years.
- Child and parents ready to participate in the study and responded to the questionnaire.

Exclusion criteria

- Child below the age of 5 years and above 15 years

- Did not respond to the questionnaire.

A predefined questionnaire, prepared in English and translated to local languages, Marathi and Kannada, was handed over to all the children attending the selected schools. Children were asked to hand over the questionnaire to their parents and return with the filled questionnaire.

The questionnaire was prepared to obtain the information related to the socio-demographic factors of the family, information related to child’s age, sex and other factors, frequency of enuresis and information related the risk factors.

The required sample size was calculated by considering the prevalence of nocturnal enuresis as 12.6 % as reported by a study conducted amongst the school going children in Lucknow.⁵ With alpha=0.05 and allowable relative error of 10 %, the calculated required sample size was 1360, to estimate the prevalence within 10 % relative error. Nocturnal enuresis was defined if the frequency of enuresis was more than twice per week 17.

Nocturnal enuresis was defined as secondary, if the child had attained full control over his nocturnal bladder habits for a continuous period of at least six months. Otherwise the nocturnal enuresis was defined as primary.

Statistical analysis

Univariate analysis was done using appropriate statistical tests like chi-square test, t test etc. Data was analysed using SPSS 23.

RESULTS

The total number of questionnaires distributed were 1710 in the selected four schools. Out of these 1710 questionnaire distributed, only 1430 completely filled questionnaire were returned.

Table 1: Age wise prevalence of nocturnal enuresis.

Age Group	Nocturnal Enuresis	Percentage	No nocturnal Enuresis	Percentage	Total
5-6	24	15.69	129	84.31	153
6-7	25	17.61	117	82.39	142
7-8	28	18.92	120	81.08	148
8-9	31	22.96	104	77.04	135
9-10	21	14.29	126	85.71	147
10-11	16	11.59	122	88.41	138
11-12	11	7.91	128	92.09	139
12-13	6	3.95	146	96.05	152
13-14	1	0.71	140	99.29	141
14-15	0	0.00	135	100.00	135
Total	163	11.40	1267	88.60	1430

($\chi^2 = 77.65$, d. f. = 7; $p = 4.15 \times 10^{-14}$)

Table 2: Sex wise prevalence of nocturnal enuresis.

Sex	Nocturnal enuresis	Percentage	No nocturnal enuresis	Percentage	Total	OR (95% CI)
Boys	105	14.34	627	85.66	732	1.85 (1.32-2.59)
Girls	58	8.31	640	91.69	698	
Total	163	11.40	1267	88.60	1430	

($\chi^2= 12.88$, d. f. = 1, p = 0.0003)

So, the response rate was 83.63%. Out of that 657 were from Marathi medium schools and 773 were from English medium schools.

The response rate from English medium schools (87.64 %) was better than the response rate from Marathi Medium schools (79.35 %). The overall prevalence of nocturnal enuresis was 11.4 %. The maximum prevalence was found in the age group of 8-9 years (22.96 %). The prevalence of nocturnal enuresis in the age group of 7 -8

was 18.92 %. There was only one child suffering from nocturnal enuresis in the age group of 13-14, whereas no child complained of it above the age of 14 years (Table 1). The differences in the prevalence rates in these age groups were statistically significant. ($\chi^2= 77.65$, p = 4.15 x 10⁻¹⁴). The prevalence of nocturnal enuresis in boys (14.34 %) was significantly more than the prevalence in girls (8.31 %). The odds ratio was 1.85 (95 % confidence interval-1.32-2.59) (Table 2). The prevalence of nocturnal enuresis was more in Marathi medium school as compared with English medium school (Table 3).

Table 3: Prevalence of nocturnal enuresis as per the medium of school.

Sex	Nocturnal enuresis	Percentage	No nocturnal enuresis	Percentage	Total	OR (95% CI)
English	65	8.41	708	91.59	773	0.52 (0.38-0.73)
Marathi	98	14.92	559	85.08	657	
Total	163	11.40	1267	88.60	1430	

($\chi^2= 14.89$, d. f. = 1, p = 0.0001)

Table 4: Association of nocturnal enuresis with various characteristics.

Characteristic	Nocturnal enuresis				Total	OR	P
	Yes	%	No	%			
Maternal education less than graduation	55	12.14	398	87.86	453	1.11 (0.79-1.57)	0.55
Paternal Education less than graduation	32	13.06	213	86.94	245	1.20 (0.80-1.83)	0.37
Below poverty line family	81	13.55	517	86.45	598	1.43 (1.03-1.99)	0.03
Stressful events	35	17.07	170	82.93	205	1.76 (1.17-2.65)	0.005
Hyperactive child	47	17.74	218	82.26	265	1.95 (1.34-2.82)	0.0003
Poor school performance	51	15.79	272	84.21	323	1.67 (1.17-2.38)	0.004
Sleep pattern (hard to awaken)	113	20.14	448	79.86	561	4.13 (2.90-5.88)	<0.0001
Family history in first blood relative	76	28.46	191	71.54	267	4.92 (3.49-6.94)	<0.0001
History of burning micturition	45	36.59	78	63.41	123	5.81 (3.85-8.78)	<0.0001
No habit of going to micturate before bed	71	19.61	291	80.39	362	2.59 (1.85-3.62)	<0.0001
Birth order (first)	79	10.84	650	89.16	729	0.89 (0.64-1.24)	0.495
Type of family (nuclear)	109	11.53	836	88.47	945	1.04 (0.74-1.47)	0.921

Table 4 shows the association of nocturnal enuresis with various socio-demographic and other factors. It shows that the nocturnal enuresis was significantly associated with socio-economic status, stress, poor school performance, sleep pattern (hard to awaken), family history, burning micturition, not going to toilet before bed. But it was not associated with maternal or paternal education, birth order, type of family. A total of 102 (62.58 %) children’s parent felt that it was an abnormal

condition and consulted any doctor. Remaining 61 parents felt that the nocturnal enuresis or bedwetting is a common and normal phenomenon.

DISCUSSION

The present cross-sectional study was conducted in school going children in the age group of 5 to 15 years to estimate the prevalence of nocturnal enuresis and its

determinants. The study has reported that the prevalence of nocturnal enuresis was 11.40%. The reported prevalence was slightly less than the reports of Srivastava et al, who reported it as 12.6%. However, it was more than the reports of De Sousa A et al, who reported it as 7.61 %.^{4,5}

The study has also revealed different prevalence rate as reported by various authors from different regions of the world. The prevalence of nocturnal enuresis in this study was lower than the reported studies from Stephanie Gonzalez Mejias and Kamleshun Ramphul (Dominican Republic, 27.9%), Aloni MN et al (Congo, 26%), Iduoriyekemwen NJ et al (Nigeria 21.3%), Bourquia A (Morocco, 35.0%), Järvelin MR (Finland 8.2%), Readett DR (Jamaica 50%).¹⁷⁻²² However, it was higher than a study from Hansakunachai T et al (Bangkok 3.9%).²³ The differences in the age groups and different socio-demographic characteristics were the main reasons in these different prevalence rates. The study has shown that the prevalence was more in the age group of 7-9 years. The findings are similar to the findings of Aljefri HM, De Sousa et al and Srivastava S et al.^{4,5,12} The study has also reported that with advancing age, the prevalence decreased. The findings of the study, boys are more affected than girls, are also similar to the findings of De Sousa et al, Mithani S et al, Ozden C et al, Pashpour et al. However, the results were not consistent with reports of Yousef KA and Aljefri HM.^{4,24,25,26}

The present study has also reported that there was statistically significant association of nocturnal enuresis with stress, family history of enuresis, lower socio-economic status. These findings are consistent with the findings of Chang JW et al and Akis N et al.^{27,28} Other risk factors, as reported by this study, such as history of burning micturition, poor school performance is also reported by Srivastava S.⁵ The study being a cross sectional study, cannot comment on the longitudinal nature of spontaneous resolution with increasing age. Also, there was lack of any clinical data. The study was based only on the questionnaire and responses from the parents.

CONCLUSION

Nevertheless, the study has concluded that the prevalence of nocturnal enuresis is 11.40 % in school going children and it is associated with age, sex, stress, family history, burning micturition, hyperactive child, poor school performance, sleep pattern (hard to awake) avoidance of going to micturate before sleep etc. It has also pointed out that the nocturnal enuresis was not associated with maternal or paternal education, birth order or type of family.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

- Butler RJ, Golding J, Northstone K. Nocturnal enuresis at 7.5 years old: Prevalence and analysis of clinical signs. *BJU Int.* 2005;96(3):404-10.
- Jiloha RC, Murthy RS. An epidemiological study of psychiatric problems in primary school children. *Child Psychiat Q.* 1981;14:1-12.
- Deivasigamani TR. Psychiatric morbidity in primary school children - an epidemiological study. *Indian J Psychiat.* 1990;32(3):235-40.
- De Sousa A, Kapoor H, Jagtap J, Sen M. Prevalence and factors affecting enuresis amongst primary school children. *Indian J Urol.* 2007;23(4):354-7.
- Srivastava S, Srivastava KL, Shingla S. Prevalence of monosymptomatic nocturnal enuresis and its correlates in school going children of Lucknow. *Indian J Pediatr.* 2013;80(6):488-91.
- Eung CK, Sihoe JD, Sit FK, Bower W, Sreedhar B, Lau J. Characteristics of primary nocturnal enuresis in adults: An epidemiological study. *BJU Int* 2004;93(3):341-5.
- Mithani S, Zaidi Z. Bed wetting in school children of Karachi. *J Pak Med Assoc.* 2005;55(1): 2-5.
- Yousef KA, Basaleem HO, bin Yahiya MT. Epidemiology of nocturnal enuresis in basic schoolchildren in Aden Governorate, Yemen. *Saudi J Kidney Dis Transpl.* 2011;22(1):167-73
- Karnicnik K, Koren A, Kos N, Marcun Varda N. Prevalence and quality of life of slovenian children with primary nocturnal enuresis. *Int J Nephrol.* 2012;2012.
- Kalo BB, Bella H. Enuresis: Prevalence and associated factors among primary school children in Saudi Arabia. *Acta Paediatr.* 1996;85(10):1217-22.
- Bower WF, Moore KH, Shepherd RB, Adams RD. The epidemiology of childhood enuresis in Australia. *Br J Urol.* 1996;78(4):602-6.
- Carman KB, Ceran O, Kaya C, Nuhoglu C, Karaman MI. Nocturnal enuresis in Turkey: Prevalence and accompanying factors in different socioeconomic environments. *Urol Int* 2008;80(4):362-6.
- Aljefri HM, Basurreh OA, Yunus F, Bawazir AA. Nocturnal enuresis among primary school children. *Saudi J Kidney Dis Transpl.* 2013;24(6):1233-41.
- Shah S, Jafri RZ, Mobin K, Mirza R, Nanji K, Jahangir F. Frequency and features of nocturnal enuresis in Pakistani children aged 5 to 16 years based on ICCS criteria: a multi-center cross-sectional study from Karachi, Pakistan. *BMC Fam Pract.* 2018;19(1):198.
- H Arnell B Bengtsson C Wassen T Emahazion G Anneren N Dahl. The genetics of primary nocturnal enuresis: inheritance and suggestion of a second major gene on chromosome 12q. *J Med Genet* 1997;34(5):360-5.
- Kawauchi A, Naitoh Y, Yoneda K, et al. Refractory enuresis related to alarm therapy. *J Pediatr Urol* 2006;2(6):579-82.

17. Hazza I, Tarawneh H. Primary nocturnal enuresis among school children in Jordan. *Saudi J Kidney Dis Transpl.* 2002;13(4):478-80.
18. Stephanie Gonzalez Mejias and Kamleshun Ramphul. Nocturnal enuresis in children from Santo Domingo, Dominican Republic: a questionnaire study of prevalence and risk factors. *BMJ Paediatr Open.* 2018;2(1):e000311.
19. Aloni MN, Ekila MB, Ekulu PM, Aloni ML, Magoga K. Nocturnal enuresis in children in kinshasa, democratic republic of congo. *Acta Paediatr.* 2012;101(10):e475-8.
20. Iduoriyekemwen NJ, Ibadin MO, Abiodun PO. Survey of childhood enuresis in the Ehor community, the EDO State, Nigeria. *Saudi J Kidney Dis Transpl* 2006;17(2):177-82.
21. Bourquia A, Chihabeddine K. Enuresis: epidemiological study in moroccan children. *Saudi J Kidney Dis Transpl* 2002;13(2):151-4.
22. Järvelin MR, Vikeväinen-Tervonen L, Moilanen I, et al. Enuresis in seven-year-old children. *Acta Paediatr Scand.* 1988;77(1):148-53.
23. Readett DR, Bamigbade T, Serjeant GR. Nocturnal enuresis in normal Jamaican children. Implications for therapy. *West Indian Med J* 1991;40(4):181-4.
24. Hansakunachai T, Ruangdaraganon N, Udomsubpayakul U, et al. Epidemiology of enuresis among school-age children in Thailand. *J Dev Behav Pediatr* 2005;26(5):356-60.
25. Mithani S, Zaidi Z. Bed wetting in school children of Karachi. *J Pak Med Assoc.* 2005;55(1):2-5.
26. Ozden C, Ozdal OL, Altinova S, Oguzulgen I, Urgancioglu G, Memis A. Prevalence and associated factors of enuresis in Turkish children. *Int Braz J Urol* 2007;33(2):216-22.
27. Pashapour N, Golmahammadlou S, Mahmoodzadeh H. Nocturnal enuresis and its treatment among primary-school children in Oromieh, Islamic Republic of Iran. *East Mediterr Health J* 2008;14(2):376-80.
28. Chang JW, Yang LY, Chin TW, Tsai HL. Clinical characteristics, nocturnal antidiuretic hormone levels, and responsiveness to DDAVP of school children with primary nocturnal enuresis. *World J Urol* 2012;30(4):567-71.
29. Akis N, Irgil E, Aytakin N. Enuresis and the effective factors: A case-control study. *Scand J Urol Nephrol.* 2002;36(3):199-203.

Cite this article as: Nakate DP, Vaidya SS, Gaikwad SY, Patil RS, Ghogare MS. Prevalence and determinants of nocturnal enuresis in school going children in Southern Maharashtra, India. *Int J Contemp Pediatr* 2019;6:564-8.